

Supplementary Table S3. Temperature and salinity associated with the maximum growth rate of the dinoflagellate *Margalefidinium polykrikoides* in laboratory conditions

Temperature (°C)	Salinity (psu)	Maximum growth rate (d ⁻¹)	Reference
25	34	0.41	Kim et al. (2004)
24	NA	0.19	Griffith and Gobler (2016)
25	30	0.35	Oh et al. (2010)
25	30	0.34	Oh et al. (2006)
25	NA	NA	Lim et al. (2021)
25	32	0.31	Lim et al. (2019)

NA, not available.

REFERENCES

- Griffith, A. W. & Gobler, C. J. 2016. Temperature controls the toxicity of the ichthyotoxic dinoflagellate *Cochlodinium polykrikoides*. Mar. Ecol. Prog. Ser. 545:63–76. doi.org/10.3354/meps11590
- Kim, D.-I., Matsuyama, Y., Nagasoe, S., et al. 2004. Effects of temperature, salinity and irradiance on the growth of the harmful red tide dinoflagellate *Cochlodinium polykrikoides* Margalef (Dinophyceae). J. Plankton Res. 26:61–66. doi.org/10.1093/plankt/fbh001
- Lee, C.-K., Park, T.-G., Park, Y.-T. & Lim, W.-A. 2013. Monitoring and trends in harmful algal blooms and red tides in Korean coastal waters, with emphasis on *Cochlodinium polykrikoides*. Harmful Algae 30(Suppl 1):S3–S14. doi.org/10.1016/j.hal.2013.10.002
- Lee, M. O., Choi, J. H. & Park, I. H. 2010. Outbreak conditions for *Cochlodinium polykrikoides* blooms in the southern coastal waters of Korea. Mar. Environ. Res. 70:227–238. doi.org/10.1016/j.marenvres.2010.05.005
- Lee, Y. S. & Lee, S. Y. 2006. Factors affecting outbreaks of *Cochlodinium polykrikoides* blooms in coastal areas of Korea. Mar. Pollut. Bull. 52:626–634. doi.org/10.1016/j.marpolbul.2005.10.015
- Lim, Y. K., Baek, S. H., Lee, M., Kim, Y. O., Choi, K.-H. & Kim, J. H. 2019. Phytoplankton composition associated with physical and chemical variables during summer in the southern sea of Korea: implication of the succession of the two toxic dinoflagellates *Cochlodinium* (a.k.a. *Margalefidinium*) *polykrikoides* and *Alexandrium affine*. J. Exp. Mar. Biol. Ecol. 516:51–66. doi.org/10.1016/j.jembe.2019.05.006
- Lim, Y. K., Park, B. S., Kim, J. H., Baek, S.-S. & Baek, S. H. 2021. Effect of marine heatwaves on bloom formation of the harmful dinoflagellate *Cochlodinium polykrikoides*: Two sides of the same coin? Harmful Algae 104:102029. doi.org/10.1016/j.hal.2021.102029
- Oh, S. J., Kim, C.-H., Kwon, H. K. & Yang, H.-S. 2010. Effects of water temperature, salinity and irradiance on the growth of harmful dinoflagellate *Cochlodinium polykrikoides* Margalef isolated from South Sea of Korea in 2008. Korean J. Fish. Aquat. Sci. 43:715–722. doi.org/10.5657/kfas.2010.43.6.715
- Oh, S. J., Yoon, Y. H., Kim, D.-I., Shimasaki, Y., Oshima, Y. & Honjo, T. 2006. Effects of light quantity and quality on the growth of the harmful Dinoflagellate, *Cochlodinium polykrikoides* Margalef (Dinophyceae). Algae 21:311–316. doi.org/10.4490/ALGAE.2006.21.3.311
- Suh, Y.-S., Kim, J.-H. & Kim, H.-G. 2000. Relationship between sea surface temperature derived from NOAA satellites and *Cochlodinium polykrikoides* red tide occurrence in Korean coastal waters. J. Environ. Sci. Int. 9:215–221.